

Pronounced surface stratification of soil phosphorus, potassium and sulfur under pastures upstream of a eutrophic wetland and estuarine system

Article

Accepted Version

Ryan, M. H., Tibbett, M., Lambers, H., Bicknell, D., Brookes, P., Barrett-Lennard, E. G., Ocampo, C. and Nicol, D. (2017) Pronounced surface stratification of soil phosphorus, potassium and sulfur under pastures upstream of a eutrophic wetland and estuarine system. *Soil Research*, 55 (7). pp. 657-669. ISSN 1838-675X doi: <https://doi.org/10.1071/SR16144> Available at <https://centaur.reading.ac.uk/73303/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1071/SR16144>

Publisher: CSIRO Publishing

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Pronounced surface stratification of soil phosphorus, potassium and sulfur under pastures upstream of an eutrophic wetland and estuarine system

Megan H. Ryan¹, Mark Tibbett², Hans Lambers¹, David Bicknell³, Phillip Brookes⁴, Ed Barrett-Lennard³, Carlos Ocampo⁵, Dion Nicol^{1, 3}

¹ School of Plant Biology, and Institute of Agriculture, The University of Western Australia, 35 Stirling Hwy, Crawley (Perth), WA 6009, Australia

² Centre for Agri-Environmental Research, School of Agriculture Policy and Development, University of Reading, Berkshire, RG6 6AR, United Kingdom

³ Department of Agriculture and Food Western Australia, 3 Baron-Hay Court, South Perth WA 6151, WA 6312, Australia

⁴ College of Environmental and Resource Sciences, Institute of Soil & Water Resources and Environmental Sciences, Zhejiang University, China

⁵ School of Civil, Environmental and Mining Engineering, The University of Western Australia, 35 Stirling Hwy, Crawley (Perth), WA 6009, Australia

Author for correspondence

Megan Ryan

School of Plant Biology, M084

The University of Western Australia, Crawley (Perth), WA, 6009

Ph +61 8 6488 2208, Fax +61 8 6488 1108

Email: megan.ryan@uwa.edu.au

26 **Summary Text for the Table of Contents.**

27 Movement of phosphorus off farms into waterways is detrimental to the health of
28 downstream aquatic systems through promoting algal blooms. We measured the
29 concentration of phosphorus, and other elements, in the top 100 mm of soil under a beef
30 farm and a dairy farm and found extremely high concentrations in the top 10 mm of soil.
31 Management practices that lower phosphorus concentrations in surface soil will likely
32 benefit health of downstream ecosystems through reducing phosphorus movement off
33 farm.

ABSTRACT

High concentrations of nutrients in surface soil present a risk of nutrient movement into waterways through surface water pathways and leaching. Phosphorus (P) is of particular concern, due to its role in aquatic system eutrophication. We measured nutrients under annual pastures on a beef farm and a dairy farm in the Peel-Harvey catchment, Western Australia. Soils were sampled in 10 mm increments to 100 mm depth in March, June and September. Plant litter contained ~300–550 mg kg⁻¹ Colwell-extractable P. Extractable soil P was strongly stratified, being ~100–225 mg kg⁻¹ (dairy) and ~50–110 mg kg⁻¹ (beef) in the top 10 mm and < 40 mg kg⁻¹ at 40–50 mm depth. Total P and extractable potassium were also highly stratified, while sulfur was less strongly stratified. Shoot nutrient concentrations indicated that nitrogen was often limiting and sulfur sometimes limiting pasture growth: concentrations of P were often much greater than required for adequate growth (>4 mg g⁻¹). We conclude that high P concentrations at the soil surface and in litter and shoots are a source of risk for movement of P from farms into waterways in the Peel-Harvey catchment.

Keywords: nutrient stratification, pH, organic matter, eutrophication, surface soil

INTRODUCTION

Eutrophication of naturally phosphorus (P)-limited waterways and standing waters due to the movement of P off farmland is a significant environmental problem on a global scale (Sharpley *et al.* 2015) and is not easily remediated (Jarvie *et al.* 2013; Rivers *et al.* 2013). Farmland may contribute P from diffuse sources, such as large areas of land with high concentrations of readily-mobile soil P due to past fertiliser applications, and from point sources, such as the slurry from intensive livestock enterprises (Weaver and Read, 1998; Rivers *et al.* 2013). Diffuse P sources present the greatest problem, as they are the most challenging to understand and amend. Worldwide, extensive research has been undertaken with the aim of reducing the contribution of diffuse P sources to eutrophication (Sharpley *et al.* 2015). However, the application of this research has proved problematic, and one of the many reasons is an incomplete understanding of the specific pathways for P movement within the farm and off the farm into waterways in the target environment (Sharpley *et al.* 2015).

In the coastal plain of the Peel-Harvey catchment in south-western Australia, forests and woodlands have been largely replaced with pastures consisting of winter-active annual species, which support beef and dairy enterprises (Ruprecht and George 1993). The area has a long-term mean annual rainfall of around 1000 mm, which is strongly winter-dominant (i.e. ~54% falls in winter and ~ 4% in summer). The area also mostly has a low surface gradient and, therefore, an extensive network of shallow open drains has been constructed to mitigate waterlogging and inundation (Cooper 1979). These open drains efficiently move surface water from upstream farms into shallow estuarine waters, saline, brackish and freshwater wetlands, and several lake systems including a ‘Wetland of International Importance’ under the Ramsar Convention on Wetlands (Hale and Butcher

2007). The movement of P into these ecosystems has long been recognised as a major contributor to eutrophication and algal blooms (Lukatelich and McComb 1986), which undesirably affect amenity, water quality and wildlife. Moreover, the Peel-Harvey catchment is part of a terrestrial biodiversity hotspot and many of its native plant species are adapted to low soil P and are poorly competitive if soil P increases (Lambers *et al.* 2013). The movement of P through overland flow or shallow groundwater flow from farms into remnant native vegetation is, therefore, also a matter of concern.

The Environmental Protection Authority of Western Australia has set a goal of reducing the movement of P into the Peel-Harvey estuary by ~50% (EPA 2008), but recent modelling suggests that P inputs from the watershed will continue to increase (Rivers *et al.* 2013). An incomplete understanding of how P moves from farms into waterways has impeded the design of management strategies to reduce P losses from soils in this catchment. Point sources of P, such as piggeries, are easily identified and relatively well understood. Diffuse sources are not as easily mapped or well understood, but are believed to contribute the most P into the estuary (Rivers *et al.* 2013). The major diffuse source of P is pastures (beef and dairy). These have been fertilised for many years, primarily with inorganic P fertiliser, and are consequently greatly enriched with P (Rivers *et al.* 2013) due to accumulation in the surface soil of P in excess to pasture requirements in both inorganic and organic forms (McLaren *et al.* 2015). The large area of pastures in the catchment accounts for most of the 2000 tonnes of P that are applied each year. Although losses of P per hectare are small they are estimated to contribute around 140 tonnes of P per year to the estuary (Rivers *et al.* 2013). Modifying farm management to reduce the movement of diffuse-source P into the open drains requires a detailed understanding of the distribution of P on these farms and the means by which P moves off-farm. Other nutrients have also

101 been supplied to pastures, notably potassium (K) and sulfur (S). Lesser enrichment with K
102 and S has likely resulted in limitations of K and S to pasture growth and P use (Weaver and
103 Reed 1998).

104
105 In the Peel-Harvey catchment, leaching of P into shallow water tables is often the primary
106 path of P movement into waterways and open drains located at low points in the landscape
107 (Weaver and Summers 2014). However, in areas with very limited slope and/or subsurface
108 impediments to groundwater flow, the movement of P through surface water pathways
109 (infiltration excess or saturation excess overland flow and, perhaps, return flow of
110 infiltrated water) may dominate, particularly once the soil profile is saturated (Ruprecht
111 and George 1993). In pastures, high concentrations of P on the soil surface due to top-
112 dressing of fertiliser, manure and infrequent tillage would increase the likelihood of P
113 movement into the open drains (Dougherty *et al.* 2006). Indeed, several studies in other
114 regions have reported high concentrations of P in the soil surface layers of pastures (0–40
115 or 0–50 mm) (McLaughlin *et al.* 1990; Haynes and Williams 1992; Cayley *et al.* 2002;
116 Dougherty *et al.* 2006) which suggests that management strategies to reduce the movement
117 of P should focus on this part of the profile. However, it is not well known whether these
118 concentrations change over time as could occur due to management such as heavy periods
119 of intensive grazing or fertiliser addition, or seasonal change such as growth of annual
120 pastures or heavy winter rainfall. Furthermore, the construction of soil P risk frameworks
121 for this catchment (e.g., soil P change point – Heckrath *et al.* 1995) would likely be
122 complicated by a heterogeneous distribution of nutrients and buffering materials through
123 the soil profile.

In view of the above, the aim of this study was to examine the stratification of P within the top 100 mm of soil in three permanent pastures in the Peel-Harvey catchment to identify whether P is highly stratified and present in high concentrations at the top of the soil profile. We also examined whether this stratification differs for pastures under contrasting management or on differing soil types and whether it changes during the course of the winter growing season. Three pastures were sampled: two mixed-composition annual pastures on a relatively low-intensity commercial beef farm (one on a deep sand and the other on a sandy loam over clay) and one annual pasture of ryegrass (*Lolium rigidum* L.) on a sand over clay on a high-intensity dairy farm (Fig. 1). The study builds upon previous studies of nutrient stratification in pastures and crops (McLaughlin *et al.* 1990; Haynes and Williams 1992; Cayley *et al.* 2002; Dougherty *et al.* 2006; Vu *et al.* 2009; Saarela and Vuorinen 2010; Haygarth *et al.* 1998) by including, in a single study, three sampling times within the growing season, small (10 mm) soil depth increments and measurements of extractable potassium and sulfur, pH and soil organic matter in addition to extractable and total P. The paddocks were under commercial management during the study.

MATERIALS AND METHODS

Environment

Pastures were sampled on two commercial farms located approximately 8 km apart (Fig. 1A), close to the town of Waroona in the coastal plain of the Peel-Harvey catchment in Western Australia. Both farms were likely established around 100 years ago following clearing of natural vegetation and soils are infrequently tilled. Fertiliser inputs across the region increased substantially after world war two (Birch, 1982; Cooper, 1979). The Peel-Harvey catchment has a typical Mediterranean climate with hot, dry summers and mild, wet winters (Table 1). Non-irrigated pastures such as the ones sampled for this study are,

therefore, largely based on winter-active annual pasture species. The two farms were chosen to be representative of two common, but contrasting, farming enterprises in the coastal plain: low-management-intensity beef enterprises and high-management-intensity dairy enterprises. Across the catchment, these two enterprise types contribute greater than 75% to fertiliser P inputs and to P entering the estuary (Rivers *et al.* 2013).

Rainfall

The monthly rainfall totals were obtained from the nearby town of Waroona (32.85°S, 115.92°E, 40 m elevation, Bureau of Meteorology station 009538) (Table 1). In the year before the study, 2011, total rainfall was similar to the long-term average. In 2012, monthly rainfall totals were close to long-term averages, except for July which received only 47 mm compared with the long-term average of 188 mm.

Beef cattle farm (two sample sites)

The beef farm had previously been a mixed-beef and dairy enterprise, but had solely been a beef-beef enterprise for more than 20 years at the time of sampling. The pasture we sampled was located on a small 3–4 m high dune of sand overlying clay, a common landform in the area. The farm was managed in a relatively low-intensity manner with an annual addition by top-dressing onto growing pastures of 100–200 kg ha⁻¹ of mineral fertiliser (7% P, 2% K, 2% S, 26% Ca, 1% Mg) and infrequent application of locally-sourced lime-sand at ~1 t ha⁻¹. No nitrogen (N) fertiliser was applied as the farmer relied upon annual legumes, primarily *Trifolium subterraneum* L., to biological fix N. The pasture contained a mix of weeds, pasture legumes and grasses, including some summer-active perennials, which were grazed by set-stocked cattle. Two sites, around 25 m apart, were sampled (Fig. 1B). The ‘upslope’ site was at the top of the dune where the soil is a

Gavin sand (McArthur and Bettenay 1959). The major pasture species present during winter were *Arctotheca calendula* (L.) Levyns, *Lolium rigidum* L., *Lupinus cosentinii* Guss., *Cynodon dactylon* (L.) Pers. and *Bromus* spp. The ‘midslope’ site was lower on the dune, close to the break of the slope towards a flat area. The soil is a Mayfield series sandy loam over clay (McArthur and Bettenay 1959) and the major pasture species present were *A. calendula*, *L. rigidum*, *Lotus subbiflorus* Lag., *T. subterraneum* and *Pennisetum clandestinum* Chiov. At the bottom of the dune on which the two sites were located is the Mayfield drain (Fig. 1B), an open drain approximately 3 m wide and 1.6 m deep. Shallow groundwater enters this drain via seepage from the bank and, more significantly, surface water enters the drain by overland flow.

Dairy cattle farm (one sample site)

The dairy farm had been under continuous management as a dairy for ~30 years. The pasture we sampled was located on a Coolup series sand over clay (~0.4 m) (Fig. 1C). It was re-sown annually with *L. rigidum* using zero-tillage techniques and contained a small amount of *Poa annua* L. and *T. michelianum* Savi. A dairy herd rotationally grazed the pasture on a tightly-controlled cycle whereby paddocks were temporarily divided to allow stock access to adequate pasture for a single day; this equated to three or four days grazing in a paddock every month. In-season N management was a priority with ~40 kg ha⁻¹ of N as urea top-dressed onto pastures after each grazing event. Lime (2.5 t ha⁻¹) was applied before the first heavy rains of the 2012 growing season on paddocks with acid soils to maintain the soil pH at 5.5. Extractable soil P concentrations were high due to past application of P fertiliser and, consequently, the farmer had not applied P fertiliser for around five years. However the farmer did apply an inorganic fertiliser containing P in 2012 (details unknown). Groundwater and, in particular, surface water from this site drain

into the South Coolup Drain (Fig. 1C), an open drain approximately 5 m wide and 1.7 m deep (Steele, 2008). Overland flow discharges via shallow-interconnected surface depressions (see arrows in Fig. 1C). More detail on drainage characteristics and nutrient export from the South Coolup Drain can be found in Steele (2008) and EPA (2008).

Sample collection

Samples were collected in March, June and September 2012. The March sampling preceded the first heavy rains of the growing season and there was little or no green pasture present. Three quadrats (0.5 × 0.5 m) were randomly placed within each of the three sites at each time of sampling; areas that had obviously recently received urine or dung were avoided. Shoots were cut close to the soil level and removed. Green shoot material in contact with the soil and senesced shoot material at the soil surface were removed (hereafter referred to as 'litter'). A custom-made 'plane' was then used to remove the soil in 10 mm increments down to 100 mm depth and samples were transported to the laboratory on the day of sampling.

Sample analyses

Soil samples were oven-dried at 40°C for one week and sieved to 2 mm. A subset of increments was analysed (i.e. 0–10, 10–20, 20–30, 30–40, 40–50, 70–80, 90–100 mm) by CSBP laboratories (Bibra Lake, Western Australia). Unless otherwise specified, the methods for soil analysis followed those of Rayment and Lyons (2011) and codes from this reference are supplied: total P (9A3b); bicarbonate-extractable P (9B) and potassium (K) (18A1, Colwell (1965)); P-buffering index (PBI) (912C, Allen and Jeffrey (1990)); extractable sulfur (S) (10D1; Blair *et al.* (1991)); pH in a soil:solution ratio of 1:5 (4A1, 4B3, 3A1); and organic carbon (6A1, Walkley and Black (1934)).

The litter samples were treated as soil samples for analysis as they were a mixture of plant material and surface soil. Samples were dried at 60°C to a constant weight. For extractable P, samples were ground to <2 mm and 1 g shaken with 100 mL of 0.5 M NaHCO₃ (pH 8.5) for 16 hours (Colwell 1963), centrifuged, and the P in the clear supernatant measured colorimetrically (Murphy and Riley 1962). For total P, samples were ground to <0.5 mm and 0.3 g digested in nitric/perchloric acid (Kuo 1996), and the P measured colorimetrically (Murphy and Riley 1962).

The pasture shoots (June and September only) and litter (March and September only) were dried at 70°C to a constant weight and then ground. Shoot samples of approximately 100 mg were digested in nitric/perchloric acid and analysed for P, K and S using inductively-coupled plasma atomic absorption with a Perkin Elmer Optima 5300 DV optical emission spectrometer (OES; Shelton, CT, USA). Nitrogen concentration was determined by dry combustion using an elemental CN analyser (Elementar Analysensysteme GmbH, Hanau, Germany).

Data analyses

Soil plane data were analysed with three-way ANOVA using Genstat version 14.1 (Lawes Agricultural Trust, Rothamsted Experimental Station, Harpenden, UK). The factors examined were site (beef midslope, beef upslope, dairy farm), depth (0–10, 10–20, 20–30, 30–40, 40–50, 70–80, 90–100 mm) and time (March, June, September) and their interactions. Most data required log₁₀ transformation to meet the assumption of normality. No outliers were removed except for the extractable P, total P and extractable K data from one quadrat at the dairy farm in September 2012 because very high concentrations at all

depth increments were interpreted as likely resulting from a feeding station previously being present (215–464 mg kg⁻¹ extractable P, 677–1333 mg kg⁻¹ total P, 152–679 mg kg⁻¹ extractable K). The litter and shoot nutrient concentration data were analysed with two-way ANOVA. The two factors examined were site (beef midslope, beef upslope, dairy farm) and time (March and September for the litter; June and September for shoots) and their interactions.

RESULTS

The results from the soil planes are presented in Figures 2–4. All variables were strongly affected by depth ($P < 0.001$), with the exception of PBI (Table 2). Sampling time strongly affected pH and extractable K ($P < 0.001$) as well as total P, S, and organic carbon ($P = 0.01$ – 0.018). Site affected all variables ($P \leq 0.004$). There were numerous interactions among factors.

Extractable P concentration was highest at the dairy site ($P < 0.001$) and declined sharply with soil depth ($P < 0.001$); indeed by 40–50 mm depth, the concentration of extractable P was generally half or less than that at 0–10 mm depth (Fig. 2A, Table 2). There was an interaction between soil depth and site ($P = 0.043$), as the decline with depth was most marked for the dairy site where the extractable P concentration at the three sampling times was ~100–230 mg kg⁻¹ at 0–10 mm, but only ~40 mg kg⁻¹ by 30–40 mm depth. There was also an interaction between site and time ($P = 0.038$) driven largely by the dairy site where extractable P concentration was lowest in June and highest in September.

The concentration of total P also declined greatly with depth ($P < 0.001$) and was affected by sampling time, being lower in March than in June and September ($P = 0.018$) (Fig. 2B,

Table 2). The concentration of total P was higher at the dairy and beef midslope sites than at the beef upslope site ($P<0.001$). At the beef upslope and dairy sites, total P was more than 200 mg kg^{-1} higher at the top of the profile in June than in March. The effect of sampling time on extractable P concentration did not always mirror total P concentration; for instance, the very high extractable P concentration in the top 10 mm of the profile in September at the dairy farm was not reflected in the total P concentration.

PBI was measured only in June and September, and was higher at the beef midslope site than at the beef upslope or dairy sites ($P=0.004$) (Fig. 2C, Table 2). PBI was unaffected by depth or sampling time ($P>0.05$). PBI was generally low (i.e. 20–50), although some samples from the beef farm were moderate (i.e. 50–100). There was large variation among the three replicate quadrats for the beef upslope and beef midslope sites, hence the three quadrats from each site are presented individually. The variation among the replicate quadrats at each site resulted from one quadrat having higher or lower PBI at all depths. The beef farm quadrats with higher PBI (i.e. >40) also had a higher total P concentration, and thus there were large standard errors for mean total P (Fig. 2B). For the beef farm, where PBI was variable, there was a significant linear positive correlation between PBI and total P concentration for the upslope and midslope sites (both $r^2=0.66$, $P<0.001$) and a significant correlation between PBI and extractable P concentration for the upslope site only ($r^2=0.32$, $P<0.001$).

The dry mass of litter on the soil surface was affected by an interaction of site and time ($P=0.006$), being similar at all three sites in March, but greatly reduced by September at the beef farm, but not the dairy (Table 3). The extractable P concentration for the litter ranged from $309\text{--}559 \text{ mg kg}^{-1}$ and was not affected by site or sampling time ($P>0.05$). The total P

concentration of the litter ranged from 525–826 mg kg⁻¹ and differed among sites ($P < 0.001$) being lowest for the beef midslope site and highest for the dairy site. There was no relationship between total P concentration in the litter and total P concentration in the top 10 mm of soil ($P > 0.05$). However, there was a positive linear relationship between total P concentration in the litter and extractable P concentration in the top 10 mm of soil ($R^2 = 0.45$, $P = 0.005$). On an area basis, the extractable and total P contained in the litter were both always less than 1.5 kg ha⁻¹ and highest in March ($P \leq 0.02$).

Soil sulfur concentration decreased strongly with depth ($P < 0.001$) and was highest at the dairy site ($P < 0.001$) (Fig. 3A; Table 2). Sulfur concentration also decreased over the sampling season ($P = 0.01$), particularly at the dairy site ($P = 0.015$). Soil extractable K concentration also decreased strongly with depth ($P < 0.001$) and decreased over the sampling season ($P < 0.001$), with the decrease from June to September greatest for the beef upslope site ($P = 0.021$). In general, soil extractable K concentration was lowest at the dairy ($P < 0.001$) (Fig. 3B). The beef farm quadrats with a high PBI also had high S and extractable K concentrations and this resulted in large standard errors at some sampling times (e.g. beef upslope for June).

Soil pH was affected by a strong three-way interaction of depth, sampling time and site ($P = 0.005$) (Fig. 4A, Table 2). Soil pH declined with sampling depth at all sites, being around one pH unit lower at 90–100 mm than at 0–10 mm. There were also strong effects of sampling time of ~0.5 to 1 pH unit, but these differed among sites and soil depths. For instance, pH was highest in September at the beef farm sites, but highest in June at the dairy and these seasonal changes occurred at all soil depths (0–100 mm) at the beef upslope and dairy sites, but not the beef midslope site. In general, soil pH was lowest at the dairy.

Soil organic carbon decreased with depth ($P < 0.001$), decreased over time ($P < 0.001$) and was lowest at the beef upslope site, intermediate at the beef midslope site and highest at the dairy ($P < 0.001$) (Fig. 4B, Table 2). An interaction among site and sampling time ($P = 0.03$) resulted from the greatest increase occurring from March to June for the beef upslope and dairy sites, but from June to September for the beef midslope. The large standard errors at the beef farm resulted from quadrats with higher organic carbon at all depths and, again, these quadrats were those with a high PBI.

Pasture shoot nutrient concentrations were measured on a bulked sample that consisted of all species present in the quadrat (Table 3). Shoot P concentration was lower for the beef midslope site than for the other two sites ($P < 0.001$) and lower in September than in June ($P = 0.002$). Shoot K concentration showed a similar trend, but this was not significant ($P > 0.05$). For shoot S concentration there was an interaction between site and sampling time ($P = 0.013$) as it did not differ among sites in June, but in September the shoot S concentrations at the beef farm sites were lower than the dairy values. Shoot N concentration was measured only in September and differed with site ($P = 0.004$) being very low for the beef farm sites ($11\text{--}13\text{ mg g}^{-1}$) compared with the dairy farm (27 mg g^{-1}).

DISCUSSION

Did P, K and S concentrations and other soil properties vary among sampling times?

For the soil nutrients we examined, the effects of sampling time were mostly small (variance ratio 2-11) compared with the effects of depth (variance ratio 14-46). Most large differences among sampling times occurred in the top 10 mm of the soil profile. Some effects of sampling time are perhaps related to the addition of fertiliser during the growing

season, likely a single application on each farm, or sampling following an intensive period of grazing (e.g., the higher extractable P and S concentrations in the surface 10 mm at the dairy farm in September). However, for extractable P, the explanation may be more complex. At the dairy farm in September, the increase in extractable P concentration of ~100 mg kg⁻¹ compared with June was far greater than could be expected from even a generous rate of fertiliser application, and was not reflected in an increase in total P. Several factors may have contributed to this increase including P mineralisation processes, perhaps associated with soil being waterlogged at the time of sampling (Bradley *et al.* 1994) and warming spring temperatures. In addition, significant mobilisation of P likely occurred during rainfall events due to leaching from the large biomass of pasture shoots that accumulated before each short grazing period. Most of the P in such green plant material is water soluble and, if leached, available to plants and soil microbes (Bromfield and Jones 1972; Noack *et al.* 2012), where some organic forms will be readily mineralised (Nash *et al.* 2014). Support for this contention comes from McDowell *et al.* (2007) who, using simulated rainfall, found that runoff from plots with membranes covering the soil (i.e. where only growing plants could contribute to nutrient load) contributed around half of the dissolved reactive P of plots where the soil interacted with the rainfall. Finally, seasonal variation in soil P pools may also result from turnover in soil microbial biomass in response to processes such as waterlogging (dairy farm) and wet/dry cycles, especially at the end of a hot, dry summer period (dairy and beef farms) (Sparling *et al.* 1985; Blackwell *et al.* 2009).

In our study, soil organic matter content tended to increase as the season progressed; this was particularly obvious for the dairy farm at 50–100 mm depth. Presumably this trend reflects the proliferation of roots of the mainly annual pasture species as the growing

season progressed as well as senescing plant material accumulating at the soil surface. Soil pH also increased by 0.5–1 units at the beef farm between March and September which was most evident in the top 0–50 mm. This increase is consistent with the return of organic matter alleviating acidity due to the growth of annual pastures and the return of senesced/trampled plant material to the soil surface.

Were P, K, S concentrations and other soil properties stratified?

At the beef and dairy farms, extractable P and total P were all highly stratified, resulting in high concentrations at the soil surface. For extractable P, similar stratification was reported for *T. subterraneum*-based pastures in south-eastern Australia (McLaughlin *et al.* 1990), pastures consisting of a mix of *Lolium perenne* L. and *T. repens* L. in New Zealand (Haynes and Williams 1992), fertilised perennial grass pastures in Finland (Saarela and Vuorinen 2010), and grass swards dominated by *L. perenne* in southwest England (Haygarth *et al.* 1998). In our study, the greatest stratification was for total P at the dairy farm where concentrations ranged from ~600–800 mg kg⁻¹ at 0–10 mm to <200 mg kg⁻¹ at 90–100 mm depth. The dairy farm had the highest concentrations of total and extractable P, reflecting a history of higher fertiliser inputs than the beef farm due to around 30 years of continuous operation as a dairy. Potassium was also highly stratified. However, S was stratified to a lesser extent, which agrees with Coad *et al.* (2010) and may reflect its greater mobility in soil (Watson 1969).

The pH of the soil decreased with depth in the top 100 mm of soil by close to one pH unit. Such pH gradients develop quite quickly under field conditions, within 5–7 years of surface soil being mixed (Conyers and Scott 1989; McLaughlin *et al.* 1990). Indeed, Evans *et al.* (1998) found that stratification developed during a single growing season, due to a

decrease in pH below 20 mm, and attributed this to nitrification followed by nitrate leaching. Organic carbon concentrations also markedly decreased with soil depth. This change was similar to that reported under crops and pastures in south-eastern Australia (McLaughlin *et al.* 1990; Vu *et al.* 2009), and is presumably largely a reflection of return of plant material to the surface of the soil.

What caused the stratification of nutrients?

The positive correlation between PBI and total P concentration at the beef farm resulted from some individual profiles having a higher PBI throughout the profile and hence enabling greater amounts of extractable P to be tightly bound. However, the very high extractable P concentrations in the soil surface layers indicates that this binding capacity had been saturated and that, consequently, P had accumulated as extractable P. Overall, the high concentrations of P and other nutrients at the soil surface likely result from the continuous input to the soil surface of plant material and animal dung (for P) and urine (for K) (Haynes and Williams 1992), as well as fertiliser (Haynes and Williams 1992; Saarela and Vuorinen 2010; McLaren *et al.* 2015, 2016). Maintenance/increase of these high surface concentrations is also exacerbated by continuous cycling of P between aboveground and belowground pools as illustrated by the strong positive linear relationship between total P concentration in the litter and extractable P concentration in the top 10 mm of soil.

Is stratification a problem?

The high concentrations of P at the soil surface undoubtedly present a significant risk for P loss during the winter wet season (Haygarth *et al.* 1998; Dougherty *et al.* 2006; Melland *et*

al. 2008) by surface water pathways (Simmonds *et al.* 2016), and also preferential flow
 pathways in the soil profile or leaching through the soil matrix (likely at the beef farm
 upslope); the latter may become prominent in the winter growing season, when surface soil
 becomes saturated and evapotranspiration is low. The very high concentrations of
 extractable P in the top 10 mm of the soil profile (i.e. up to $\sim 225 \text{ mg kg}^{-1}$) may exacerbate
 P movement if they are above the “change point” (Heckrath *et al.* 1995). The change point
 has not been determined for these soils. However, key challenges for using it, or other soil
 character-defined parameters, to define the risk of P movement off pastures in the Peel
 Harvey arise from the high P stratification in the top 100 mm of the soils and the variation
 in PBI over small scales. As mixing contrasting samples from depth increments or
 replicate profiles likely substantially changes results, we suggest that further work is
 required to adapt risk prediction for stratified, variable soils.

How can stratification be reduced?

Several management practices have been proposed as a means to reduce P loss from farms
 into waterways in the Peel-Harvey catchment. These include using perennial pastures,
 managing waterways and riparian vegetation, only applying P fertiliser if soil tests indicate
 P is limiting for pasture growth, using less-soluble fertilisers, applying fertiliser onto
 growing pastures and using P-retentive soil amendments (Rivers *et al.* 2013). Our finding
 of a high surface concentration of nutrients suggests that other management practice
 changes should be considered.

Selection of more P-efficient pasture cultivars is one possible change. Species and cultivars
 may differ greatly in shoot P concentrations at a given level of extractable soil P, and in
 external critical P requirements, with grasses generally presenting lower shoot P

concentrations and a lower external critical P requirement than legumes (Ozanne *et al.* 1969; McDowell *et al.* 2011; Sandral *et al.* 2015; Haling *et al.* 2016). Thus, decreasing the rate of P-fertiliser addition to pastures to match the P requirements of grasses only, or of annual legumes with lower P requirements than the current widely-used species *T. subterraneum*, would reduce the size of the pool of readily-soluble P circulating through plants and top soil layers and thus reduce the risk of losses. As applying P fertiliser above the rates required for pasture maintenance causes accumulation of sparingly-soluble forms of organic and inorganic P in the soil (McLaren *et al.* 2015), reduced rates of P fertiliser application would also improve the P use efficiency of the farming systems.

Mixing of soil through tillage or soil inversion may reduce the risk of nutrient loss from top soil layers through surficial processes by reducing nutrient concentrations at the soil surface (Vu *et al.* 2009) and in the soil water at the surface (Nash *et al.* 2015). Soil amendments to aid P retention could also be incorporated at the time of tillage (e.g., Summers *et al.* 1996). However, the impact of mixing soil high in P lower into the profile on the loss of P into shallow water tables may require consideration in some instances (e.g. the dairy farm in our study). Interestingly, soil inversion may have desirable side-effects such as those shown for ‘clay delving’ where bringing subsoil clay to the surface improves profile wettability and reduces preferential flow (Betti *et al.* 2015). Indeed, this sort of ‘strategic tillage’ is increasingly considered as beneficial for minimum tillage cropping systems (Dang *et al.* 2015), which also accumulate P in the top 5 mm of soil (Vu *et al.* 2009). However, the reduction in nutrient stratification from tillage or soil inversion will dissipate over time (Nash *et al.* 2015). For instance, McLaughlin *et al.* (1990) rotary-hoed the top 100 mm of soil under an annual pasture, applied 250 kg ha⁻¹ year⁻¹ of superphosphate, and then cut and removed plant shoots annually. After seven years,

stratification was similar to that in pastures that were not rotary hoed. Hence, in the Peel-Harvey coastal catchment, the ideal frequency of tillage or soil inversion conducted to reduce nutrient stratification would need to be determined.

Limitations to pasture growth

Whilst interpretation of our shoot nutrient concentration data must be undertaken with care as the botanical composition of the pastures varied, they do suggest S and N limitations to pasture growth at the beef farm in September and an N limitation at the dairy farm (Weir and Cresswell 1994). This is consistent with the findings of Weaver and Reed (1998) who sampled soils across the south coast region of Western Australia finding that two thirds of high P status soils were deficient in S and a quarter deficient in K. Note that the proliferation of kikuyu (*P. clandestinum* Chiov.) may explain the low shoot P concentrations at the beef midslope site (Fulkerson *et al.* 1998). Limitations to pasture growth from S, K and N, along with the over-application of P fertiliser, may have been the cause of the high concentrations of P ($>4 \text{ mg g}^{-1}$) in pasture shoots at the dairy and beef upslope sites, and created a large pool of readily-leachable P in the living pasture shoots (Bromfield and Jones 1972; McDowell *et al.* 2007). Addressing these nutrient limitations may result in dilution of shoot P concentrations and potentially reduce the proportion of shoot P consisting of reactive/soluble P; it would also improve pasture production.

CONCLUSIONS

We found pronounced stratification of P, K and S in the top 100 mm of the soil profile and very high concentrations of P at the soil surface, particularly at the dairy. The litter on top of the soil surface and pasture shoots also had high P concentrations and this was most pronounced at the dairy. Differences among the three sampling times in concentrations of

P, K and S in soil were generally small and largest in the surface layers. Some large fluctuations in extractable soil P concentration at the dairy could not be readily explained. Thus, it seems that there are gaps in our understanding of P cycling in these pastures and, particularly, of the role of plants in P cycling. Nevertheless, together, the high concentrations of P at the soil surface and in the litter and pasture shoots undoubtedly constitute a source of risk for the movement of P off farmland. While there are many management practices that can be, or have already been, adopted to reduce P movement our results suggest three areas that merit further investigation: (1) use of pasture legumes that require less P for maximum yield and have lower shoot P concentrations than current cultivars; (2) soil tillage or inversion to reduce P stratification; and (3) capacity to reduce shoot P concentrations through reduced P-fertiliser application coupled with removal of K, S and N limitations to pasture growth.

ACKNOWLEDGEMENTS

This study was funded by ARC linkage project LP120200808 in collaboration with Greening Australia, the Harvey River Restoration Taskforce, Alcoa Farmlands, the Department of Food and Agriculture Western Australia and the National Measurement Institute. Seed funding was provided by the Alcoa Foundation grant “Towards Sustainable Landscapes: Phosphorus Management to Protect Ramsar Wetlands”. Megan Ryan is funded by an ARC Future Fellowship (FT140100103). The authors also thank Neil Bruce and the late Kevin Dilley who generously allowed sampling on their farms, as well as generous collaborators Jane Townsend and Tony Hiscock.

REFERENCES

- 524 Allen DG, Jeffrey RC (1990) 'Methods of analysis of phosphorus in Western Australian
525 soils.' (Chemistry Centre of WA: Perth)
- 526 Betti G, Grant C, Churchman G, Murray R (2015) Increased profile wettability in texture-
527 contrast soils from clay delving: case studies in South Australia. *Soil Research* **53**, 125–
528 136.
- 529 Birch PB (1982) Phosphorus export from coastal plain drainage into the Peel-Harvey
530 estuarine system of Western Australia. *Australian Journal of Marine and Freshwater*
531 *Research* **33**, 23–32.
- 532 Bureau of Meteorology (2016) Climate data online. <http://www.bom.gov.au/> (accessed 1
533 August 2015)
- 534 Blackwell MSA, Brookes PC, de la Fuente-Martinez N, Murray PJ, Snars KE, Williams
535 JK, Haygarth PM (2009) Effects of soil drying and rate of re-wetting on concentrations and
536 forms of phosphorus in leachate. *Biology and Fertility of Soils* **45**, 635–643.
- 537 Bradley J, Vimpany I, Nicholls PJ (1984) Effects of waterlogging and subsequent drainage
538 of a pasture soil on phosphate sorption, extractable phosphate and oxalate-extractable iron.
539 *Soil Research* **22**, 455–461.
- 540 Blair GJ, Chinoim N, Lefroy RDB, Anderson GC, Crocker GJ (1991) A soil sulfur test for
541 pastures and crops. *Australian Journal of Soil Research* **29**, 619–626.
- 542 Bromfield SM, Jones OL (1972) Initial leaching of hayed-off pasture plants in relation to
543 cycling of phosphorus. *Australian Journal of Agricultural Research* **23**, 811–824.

- 544 Cayley JWD, McCaskill MR, Kearney GA (2002) Available phosphorus, sulfur,
545 potassium, and other cations in a long-term grazing experiment in south-western Victoria.
546 *Australian Journal of Agricultural Research* **53**, 1349–1360.
- 547 Coad JR, Burkitt LL, Gourley CJP (2010) Influence of sample depth on extractable
548 nutrient concentrations, pH and the phosphorus buffering index of pasture soils in south-
549 eastern Australia. *Australian Journal of Soil Research* **48**, 355–360.
- 550 Colwell JD (1963) The estimation of the phosphorus fertiliser requirements of wheat in
551 southern NSW by soil analysis. *Australian Journal of Experimental Agriculture and*
552 *Animal Husbandry* **3**, 190–197.
- 553 Colwell JD (1965) An automatic procedure for the determination of phosphorus in sodium
554 hydrogen carbonate extracts of soils. *Chemistry and Industry* **10**, 893–895.
- 555 Conyers MK, Scott BJ (1989) The influence of surface incorporated lime on subsurface
556 soil acidity. *Australian Journal of Experimental Agriculture* **29**, 201–207.
- 557 Cooper WS (1979) Drainage and irrigation. In 'Western landscapes'. (Ed. J Gentilli) pp.
558 245-252. (UWA Press: Nedlands)
- 559 Dang YP, Seymour NP, Walker SR, Bell MJ, Freebairn DM (2015) Strategic tillage in no-
560 till farming systems in Australia's northern grains-growing regions: I. Drivers and
561 implementation. *Soil and Tillage Research* **152**, 104–114.
- 562 Dougherty WJ, Nash DM, Chittleborough DJ, Cox JW, Fleming NK (2006) Stratification,
563 forms, and mobility of phosphorus in the topsoil of a Chromosol used for dairying.
564 *Australian Journal of Soil Research* **44**, 277–284.

- 565 EPA (2008) 'Water quality improvement plan for the rivers and estuary of the Peel-Harvey
566 system — phosphorus management.' (Environmental Protection Authority: Perth, Western
567 Australia)
- 568 Evans CM, Conyers MK, Black AS, Poile GJ (1998) Effect of ammonium, organic
569 amendments, and plant growth on soil pH stratification. *Australian Journal of Soil*
570 *Research* **36**, 641–653.
- 571 Fulkerson WJ, Slack K, Hennessy DW, Hough GM (1998) Nutrients in ryegrass (*Lolium*
572 spp.), white clover (*Trifolium repens*) and kikuyu (*Pennisetum clandestinum*) pastures in
573 relation to season and stage of regrowth in a subtropical environment. *Australian Journal*
574 *of Experimental Agriculture* **38**, 227–240. Hale J, Butcher R (2007) Ecological character
575 description of the Peel-Yalgorup Ramsar Site, Report to the Department of Environment
576 and Conservation and the Peel-Harvey Catchment Council, Perth, Western Australia.
- 577 Haling RE, Yang Z, Shadwell N, Culvenor RA, Stefanski A, Ryan MH, Sandral GA, Kidd
578 DR, Lambers H, Simpson RJ (2016) Growth and root dry matter allocation by pasture
579 legumes and a grass with contrasting external critical phosphorus requirements. *Plant and*
580 *Soil*, doi: 10.1007/s11104-016-2808-2
- 581 Haygarth PM, Hepworth L, Jarvis SC (1998) Forms of phosphorus transfer in hydrological
582 pathways from soil under grazed grassland. *European Journal of Soil Science* **49**, 65–72.
- 583 Haynes RJ, Williams PH (1992) Long-term effect of superphosphate on accumulation of
584 soil-phosphorus and exchangeable cations on a grazed, irrigated pasture site *Plant and Soil*
585 **142**, 123–133.

- 586 Heckrath G, Brookes PC, Poulton PR, Goulding KWT (1995) Phosphorus leaching from
587 soils containing different phosphorus concentrations in the Broadbalk Experiment. *Journal*
588 *of Environmental Quality* **24**, 904–910.
- 589 Jarvie HP, Sharpley AN, Withers PJA, Scott JT, Haggard BE, Neal C (2013) Phosphorus
590 mitigation to control river eutrophication: murky waters, inconvenient truths, and
591 "postnormal" science. *Journal of Environmental Quality* **42**, 295–304.
- 592 Kuo S (1996) Phosphorus. In 'Methods of Soil Analysis. Part 3 Chemical Methods'. (Eds D
593 Sparks, A Page, P Helmke, R Loeppert) pp. 869–920. (SSSA Book Series No. 5, Soil
594 Science Society of America, Inc.: Madison, WI)
- 595 Lambers H, Ahmedi I, Berkowitz O, Dunne C, Finnegan PM, Hardy GESJ, Jost R,
596 Laliberté E, Pearse SJ, Teste FP (2013) Phosphorus nutrition of phosphorus-sensitive
597 Australian native plants: threats to plant communities in a global biodiversity hotspot.
598 *Conservation Physiology* **1**, cot010.
- 599 Lukatelich RJ, McComb AJ (1986) Nutrient levels and the development of diatom and
600 blue-green-algal blooms in a shallow Australian estuary. *Journal of Plankton Research* **8**,
601 597–618.
- 602 McArthur W, Bettenay E (1959) 'The soils and irrigation potential of the Pinjarra-Waroona
603 area, Western Australia. Australia.' (CSIRO, Division of Soils: Melbourne)
- 604 McDowell RW, Nash DM, Robertson F (2007) Sources of phosphorus lost from a grazed
605 pasture receiving simulated rainfall. *Journal of Environmental Quality* **36**, 1281–1288.
- 606 McDowell RW, Sharpley AN, Crush JR, Simmons T (2011) Phosphorus in pasture plants:
607 potential implications for phosphorus loss in surface runoff. *Plant and Soil* **345**, 23–35.

- 608 McLaren TI, Simpson RJ, McLaughlin MJ, Smernik RJ, McBeath TM, Guppy CN,
 609 Richardson AE (2015) An assessment of various measures of soil phosphorus and the net
 610 accumulation of phosphorus in fertilized soils under pasture. *Journal of Plant Nutrition*
 611 *and Soil Science* **178**, 543–554.
- 612 McLaren TI, McLaughlin MJ, McBeath TM, Simpson RJ, Smernik RJ, Guppy CN,
 613 Richardson AE (2016) The fate of fertiliser P in soil under pasture and uptake by
 614 subterranean clover – a field study using ³³P-labelled single superphosphate. *Plant and*
 615 *Soil* **401**, 23–38.
- 616 McLaughlin MJ, Baker TG, James TR, Rundle JA (1990) Distribution and forms of
 617 phosphorus and aluminum in acidic topsoils under pastures in south-eastern Australia.
 618 *Australian Journal of Soil Research* **28**, 371–385.
- 619 Melland AR, Mc Caskill MR, White RE, Chapman DF (2008) Loss of phosphorus and
 620 nitrogen in runoff and subsurface drainage from high and low input pastures grazed by
 621 sheep in southern Australia. *Australian Journal of Soil Research* **46**, 161–172. Murphy J,
 622 Riley JP (1962) A modified single solution method for the determination of phosphate in
 623 natural waters. *Analytical Chemistry Acta* **27**, 31–60.
- 624 Nash DM, Watkins M, Heaven MW, Hannah M, Robertson F, McDowell R (2015) Effects
 625 of cultivation on soil and soil water under different fertiliser regimes. *Soil and Tillage*
 626 *Research* **145**, 37–46.
- 627 Nash DM, Haygarth PM, Turner BL, Condon LM, McDowell RW, Richardson AE,
 628 Watkins M, Heaven MW. (2014) Using organic phosphorus to sustain pasture
 629 productivity: a perspective. *Geoderma*. **221**, 11-19.

- 630 Noack SR, McLaughlin MJ, Smernik RJ, McBeath TM, Armstrong RD (2012) Crop
631 residue phosphorus: speciation and potential bio-availability. *Plant and Soil* **359**, 375–385.
- 632 Ozanne PG, Keay J, Biddiscombe EF (1969) Comparative applied phosphate requirements
633 of eight annual pasture species. *Australian Journal of Agricultural Research* **20**, 809–818.
- 634 Rayment GE, Lyons D (2011) 'Soil Chemical Methods – Australasia.' (CSIRO Publishing
635 Melbourne, Australia)
- 636 Rivers MR, Weaver DM, Smettem KRJ, Davies PM (2013) Estimating farm to catchment
637 nutrient fluxes using dynamic simulation modelling – Can agri-environmental BMPs really
638 do the job? *Journal of Environmental Management* **130**, 313–323.
- 639 Ruprecht JK, George PR (1993) Hydrology of the Peel-Harvey estuary catchment.
640 *Fertilizer Research* **36**, 127–133.
- 641 Saarela I, Vuorinen M (2010) Stratification of soil phosphorus, pH, and macro-cations
642 under intensively cropped grass ley. *Nutrient Cycling in Agroecosystems* **86**: 367–381.
- 643 Sandral G, Simpson R, Yang Z, Culvenor R, Kidd D, Lambers H, Ryan M (2015)
644 Phosphorus efficient pastures: response of alternative legumes to fertiliser application. In
645 'Proceeding of the 17th Australian Agronomy Conference'. (Eds T Acuña, C Moeller, D
646 Parsons, M. Harrison) (Australian Society of Agronomy: Hobart)
- 647 Sharpley AN, Bergstrom L, Aronsson H, Bechmann M, Bolster CH, Borling K, Djodjic F,
648 Jarvie HP, Schoumans OF, Stamm C, Tonderski KS, Ulen B, Uusitalo R, Withers PJA
649 (2015) Future agriculture with minimized phosphorus losses to waters: Research needs and
650 direction. *Ambio* **44**, S163–S179.

- 651 Simmonds B, McDowell RW & Condon LM (2016) The effect of soil moisture extremes
 652 on the pathways and forms of phosphorus lost in runoff from two contrasting soil
 653 types. *Soil Research*, **In Press**
- 654 Sparling GP, Whale KN, Ramsay AJ (1985) Quantifying the contribution from the soil
 655 microbial biomass to the extractable P-levels of fresh and air-dried soils. *Australian*
 656 *Journal of Soil Research* **23**, 613–621.
- 657 Steele J (2008) Management of diffuse water quality pollution in the Peel-Harvey coastal
 658 drainage system: a strategic approach to implementation of best management practices.
 659 Report prepared for Peel-Harvey Catchment Council Inc. (Mandurah), 202 p.
- 660 Summers RN, Smirk DD, Karafilis D (1996) Phosphorus retention and leachates from
 661 sandy soil amended with bauxite residue (red mud). *Australian Journal of Soil Research*
 662 **34**, 555–567.
- 663 Vu DT, Tang C, Armstrong RD (2009) Tillage system affects phosphorus form and depth
 664 distribution in three contrasting Victorian soils. *Soil Research* **47**: 33–45.
- 665 Walkley A, Black IA (1934) An examination of the Degtjareff method for determining soil
 666 organic matter, and a proposed modification of the chromic acid titration method. *Soil*
 667 *Science* **37**, 29–38.
- 668 Watson ER (1969) The influence of subterranean clover pastures on soil fertility III. Effect
 669 of applied phosphorus and sulphur *Australian Journal of Agricultural Research* **20**, 447–
 670 456.
- 671 Weaver D, Summers R (2014) Fit-for-purpose phosphorus management: do riparian
 672 buffers qualify in catchments with sandy soils? *Environmental Monitoring and Assessment*
 673 **186**, 2867–2884.

- 674 Weaver DM and Reed AEG (1998) Patterns of nutrient status and fertiliser practice on
675 soils of the south coast of Western Australia. *Agriculture, Ecosystems & Environment* **67**,
676 37–53.
- 677 Weir RG, Cresswell GC (1994) 'Plant nutrient disorders 4: Pastures and field crops.'
678 (Inkata: Melbourne)

679 Table 1. Monthly rainfall totals in 2012 and the long-term average (LTA) monthly rainfall totals and monthly maximum and minimum
 680 temperatures at Waroona, the closest town to the farms (Bureau of Meteorology, 2016). Months when sampling occurred are
 681 highlighted. *=missing data

		J	F	M	A	M	J	J	A	S	O	N	D	Total
Rainfall (mm)	2012	2	5	0	45	94	214	47	117	156	37	75	*	794
	LTA	12	14	21	54	140	205	188	149	92	57	37	14	992
Maximum temperature (°C)	LTA	30	30	27	23	19	16	15	16	17	20	24	27	
Minimum temperature (°C)	LTA	14	15	13	10	8	7	6	6	6	8	10	13	

682

683 Table 2. Outcomes of three-way ANOVAs on extractable and total phosphorus (P), P-buffering index (PBI), sulfur (S), extractable potassium
 684 (K), pH and organic carbon; the factors were depth (0–10, 10–20, 20–30, 30–40, 40–50, 70–80, 90–100 mm), sampling time (March, June,
 685 September) and site (beef midslope, beef upslope, dairy farm) and their interactions.

	Depth (D)	Sampling time (T)	Site (S)	D x S	T x S	D x T	D x T x S
Log ₁₀ Extractable P	v.r.=34, P<0.001	n.s.	v.r. = 17, P<0.001	v.r.=2, P=0.043	v.r.=3, P=0.038	n.s.	n.s.
Log ₁₀ Total P	v.r.=30, P<0.001	v.r.=4, P=0.018	v.r. = 7, P=0.001	n.s.	n.s.	n.s.	n.s.
Log ₁₀ PBI ¹	n.s.	n.s.	v.r. = 6, P=0.004	n.s.	n.s.	n.s.	n.s.
Log ₁₀ S	v.r.=14, P<0.001	v.r.=5, P=0.01	v.r. = 75, P<0.001	n.s.	v.r.=3, P=0.015	n.s.	n.s.
Log ₁₀ Extractable K	v.r.=46, P<0.001	v.r.=11, P<0.001	v.r. = 30, P<0.001	n.s.	v.r.=3, P=0.021	n.s.	n.s.
Antilog pH	v.r.=17, P<0.001	v.r.=47, P<0.001	v.r. = 36, P<0.001	v.r.=3, P<0.001	v.r.=26, P<0.001	v.r.=3, P<0.001	v.r.=2, P=0.005
Log ₁₀ Organic carbon	v.r.=8, P<0.001	v.r.=4, P=0.013	v.r. = 20, P<0.001	n.s.	v.r.=3, P=0.03	n.s.	n.s.

686 n.s., not significant, v.r., variance ratio

687

Phosphorus: Deficient - <2.0 ; Low - 2.2-2.3; Normal - 2.5-0.5 mg g⁻¹. Potassium: Normal 11-25 5 mg g⁻¹. Sulfur: Deficient - <2.2 ;
Low - 2.2-2.3 mg g⁻¹; Normal - 2.5-0.4 mg g⁻¹; Nitrogen; no values given for deficient, Low 30-32, Normal 33-55 mg g⁻¹ (Weir and
Cresswell (1994) for subterranean clover)

694 n.s., not significant, v.r., variance ratio

Figure captions


Figure 1. The location of the beef and dairy farms in the Peel Harvey catchment (A) and the location of the two sites at the beef farm (B) and the single site at the dairy farm (C). Arrows in (B) and (C) indicate surface run-off direction following topographic slope.

Figure 2. The concentration of extractable phosphorus (P) (A) and total P (B), and the P-buffering index (PBI) (C) in 10 mm increments of soil profiles sampled to 100 mm depth for three sites (mean \pm s.e., n=3). Note that for C), the three individual replicate quadrats are shown for the beef farm sites to illustrate the variability among replicates.


Figure 3. The concentration of sulfur (S) (A) and extractable potassium (K) (B) in 10 mm increments of soil profiles sampled to 100 mm depth for three sites (mean \pm s.e., n=3).

Figure 4. The pH in CaCl₂ (A) and organic carbon concentration (B) in 10 mm increments of soil profiles sampled to 100 mm depth for three sites (mean \pm s.e., n=3).

Fig 1




724 Fig 2 revised




725

726 Fig 3 REVISED



727

728 Fig 4 REVISED



729